

First insight into freshwater fish assemblages in the western part of the Endau-Rompin landscape, Malaysia

Munian Kaviarasu^{1,2}, Farah Farhana Ramli¹,
Lokman Mohd Ilham Norhakim^{1,3}, Nursyuhada Othman¹,
Nur Aina Amira Mahyuddin³, Hidayah Haris¹, Nur Hartini Sariyati¹,
Mohd Faudzir Najmuddin¹, Salim Aman⁴, Salman Faris Zaharin⁵,
Muhammad Abu Bakar Abdul-Latiff¹

1 Environmental Management and Conservation Research Unit (eNCORe), Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus), 84600 Muar, Johor, Malaysia **2** Zoology Branch, Forest Biodiversity Division, Forest Research Institute Malaysia (FRIM), 52109 Kepong, Selangor, Malaysia **3** Kim Ichthyologist Centre, Kg Parit Samsu, Jalan Temenggong Ahmad, 84150 Parit Jawa, Muar Johor, Malaysia **4** Forestry Department of Johor State, Second floor, Bangunan Dato' Mohamad Ibrahim Munsyi, 79660 Nusajaya, Johor, Malaysia **5** Department of Fishery District Batu Pahat, No 1444, Jalan Pantai, 83000 Batu Pahat, Malaysia

Corresponding authors: Munian Kaviarasu (kaviarasu@frim.gov.my)

Muhammad Abu Bakar Abdul-Latiff (latiff@uthm.edu.my)

Academic editor: Valter Azevedo-Santos | Received 20 May 2022 | Accepted 17 November 2022 | Published 16 December 2022

<https://zoobank.org/7FFB5496-4393-405C-B544-7611FC9D1865>

Citation: Kaviarasu M, Ramli FF, Mohd Ilham Norhakim L, Othman N, Mahyuddin NAA, Haris H, Sariyati NH, Najmuddin MF, Aman S, Zaharin SF, Abdul-Latiff MAB (2022) First insight into freshwater fish assemblages in the western part of the Endau-Rompin landscape, Malaysia. *Nature Conservation* 50: 265–281. <https://doi.org/10.3897/natureconservation.50.86090>

Abstract

In Malaysia, our knowledge of freshwater ecosystem and its aquatic inhabitants, particularly freshwater fish, remains inadequate, even in protected regions. It is essential to understand the composition of freshwater fish, their distribution along river gradients, and their interactions between environmental variables to develop and strategize effective conservation and management plans. Consequently, an investigation into freshwater fish assemblages in three rivers draining off from the western part of Endau-Rompin Landscape was conducted in 12 established substations. Sampling sessions were conducted from September to December 2021 in the Labis district of Johor, using multiple sampling methods along 200 meters for each substation. The environmental variables were measured using water quality parameters. A total of 66 species were collected. The family Cyprinidae presented the highest species diversity (17 species), constituting 52% of total capture. *Crossocheilus obscurus* was the most dominant species, and the highest species richness was

recorded in the Segamat River (45 species), followed by Juaseh (36 species) and Labis River (34 species). It was discerned that the composition of fish varied between the substations. The knowledge presented here is the first documentation on the freshwater fish from these rivers. It would serve as a baseline information for key authorities and stakeholders to conserve the biodiversity inhabiting freshwater ecosystems in Malaysia.

Keywords

conservation, diversity, ichthyology, redundancy analysis, species richness

Introduction

The surface freshwater habitats contain only approximately 0.01% of the world's water volume and cover only roughly 0.8% of the Earth's surface (Gleick 1996; Reid et al. 2019; Miller 2021; Val et al. 2022). Yet, it is home to a vast number of species and broadly used by human for various ecosystem services. The decline of freshwater biodiversity is far greater than in most affected other terrestrial ecosystems (Sala et al. 2000; Darwall et al. 2018; Harrison et al. 2018; Reid et al. 2019; Tickner et al. 2020). Currently, the rate of wetland loss is three times that of forest loss (Gardner and Finlayson 2018), and populations of freshwater vertebrate species have declined at twice the rate of that involving land or ocean vertebrates (Grooten and Almond 2018). Reid et al. (2019) recognized that threats, such as water pollution, overexploitation, flow modification, destruction or degradation of habitat, and invasion by exotic species persist to the present, as reviewed by Dudgeon et al. (2006) a decade ago with 12 more stressors that implicate the freshwater biodiversity globally. Similar threats to the freshwater ecosystems were reported in Malaysia and these will assuredly endanger the freshwater ecosystem and biodiversity.

Rivers in Malaysia are rich ecosystems with wider functions, including providing water supply, irrigation for agriculture, a means of transportation, a source of food (fisheries), hydroelectric power, and water use for industries (Chan 2002). Rivers are also habitats for riverine and aquatic flora and fauna, and the riparian environment supports a rich biodiversity of life forms (Naiman and Bilby 1998). However, over the years, the water quality of rivers has been substantially damaged due to a combination of factors, namely the low priority accorded to it by the government, public apathy, negligence, and bad management. (Chan 2012). With the rapid destruction of natural habitats along the freshwater ecosystem, it is essential to document the presence, diversity, and distribution of fauna, especially the freshwater fish in various inland water bodies in Malaysia.

Fishes are important ecological components of the freshwater ecosystem, and they are one of the animals currently threatened in Malaysia. According to Chong et al. (2000) habitat deterioration, loss, and change threaten almost 76% of freshwater fish species in Malaysia. Furthermore, Malaysian freshwater fish documentation remains in its early stages, and no centralized taxonomy oversight exists (Ng et al. 2017; Ng et al. 2019). Baseline information on the diversity and distribution of the species of concern is essential for developing holistic conservation and management strategies. In fact, knowledge of freshwater fish present across protected areas and surrounding ecosystems in Malaysia remains poor and inadequate.

Endau-Rompin Landscape (**ERL**) comprises two major protected forest lands; Rompin State Park (**RSP**) which is located in the State of Pahang while Endau-Rompin Johor National Park (**ERJNP**) of Johor is administered by the Johor National Parks Corporation. The ERJNP is surrounded by several permanent forest reserves including the Labis, Mersing, Kluang, Lenggong and Ulu Sedili Permanent Reserved Forests and managed by Johor Forestry Department. The ERL itself encompassed 3,600 km² of total coverage.

The ERL is located within a matrix of other land cover types, especially oil palm and rubber plantations to the north, west, and south (Saaban et al. 2020). The landscape was well-chronicled for its large mammals, including tigers and elephants, and the conflicts triggered by these large animals. Studies on other fauna groups were also well-established, especially within RPS and ERJNP. While the gazettement safeguarded the RPS and ERJNP, the forest reserves surrounding the protected forest land faced an onslaught of threats ranging from land-use conversion to agricultural and infrastructural development (Aqilah et al. 2018). It is time to initiate a comprehensive study on the ecosystem and diversity found within these forest reserves, particularly the freshwater fish assemblages that remain unassessed.

The main aim of this study was to document the freshwater fish assemblages in three rivers that flow off from the Labis Forest Reserve, a forest complex next to ERL. Second, we aim to verify if the composition of fish changes gradually along the river gradient and if the fish composition differs between the substations as well. This is the first study on freshwater fish assemblages from rivers flowing off from the western part of ERL. It is expected that the information would serve as a baseline knowledge for key authorities and stakeholders to conserve the biodiversity inhabiting freshwater ecosystems in Malaysia, and to promote conservation and management programs in the future.

Methods and materials

Study sites

Three rivers, Juaseh, Labis, and Segamat rivers, flowing off from the western part of the ERL, were assessed for their freshwater fish assemblages (Fig. 1). The average width and depth of Juaseh River was 8.51 meters and 0.4 meters respectively. It is estimated that the Juaseh River runs along approximately 29 km before joining the Segamat River. Similarly, the Segamat River with an average of 11.06 meters of width and 0.4 meters depth, originated from the ERJNP, flows approximately 45 km before joining with the Muar River at the west of the national park. For the Labis River, the dimensions are an average of 5.33 meters in width and depth ranged between 0.15–0.20 meters. The river starts from the hills of the Labis Forest Reserve, then flows to the south of the landscape complex before joining the Muar River. It is estimated that the Labis River runs more than 60-km before flowing into the Muar River.

Four substations were established in each river with a minimum of 3 km intervals between one substation and another. Each substation sampled its freshwater fish along a 200-meter long transect. All substations #1 were located at the headwater of each




Figure 1. The localities of the 12 substations to study the composition and distribution between freshwater fish assemblages in three rivers in ERL.

river, where they were protected in the forest reserve, and the river's condition was pristine from any anthropogenic activities. The vegetation conditions in each substation #1 were dense with emerging trees, and the rivers were covered with close canopies. The soil substrates were mostly boulders with sandy banks. Some sites were designated as public recreation sites, such as waterfalls.

The subsequent substations #2, #3, and #4 of each river flowed through developed areas, including open land, shrubs, housing areas, and oil palm plantations. Largely, this section of the river was lightly polluted thanks to the presence of oil palm plantations or human settlements. The vegetation along the substation was mainly shrubs with algae submerged into riverbeds and was not covered by tree canopies. The boulders and cobbles were absent in these substations but were largely covered by sand, silt, and clay (Table 1).

Freshwater fish sampling

Fish samples were collected from 12 substations (Fig. 1) based on habitat representativeness and accessibility from September to November in the year 2021. Fish sampling in each substation was based on two approaches; electro-fishing and net scooping along the 200-m transect. The first group with a minimum of four persons were tasked with electro-fishing, while the second group conducted scooping using fine-sized scoop nets along the transect. A backpack Root-Smith LB24 electro-fisher unit was used to stun fish in each designated substation by applying a standardized voltage and current for an hour of effort. A single person operated the unit with assistance from three to four persons to collect the stunned fish using scoop nets. The second group was tasked with scooping out fish using fine-sized scoop nets from underneath wood debris, below riverbanks with hollows, sub-

Table 1. The locations and characteristics of substations established at the western part of ERL, Malaysia.

River	GPS coordinates	Width (m)	Depth (m)
Segamat River			
Substation #1	2°31'3.10"N, 103°6'25.91"E	14.57	0.51
Substation #2	2°32'45.56"N, 103°0'42.09"E	10.1	0.36
Substation #3	2°33'4.29"N, 102°59'18.55"E	7.15	0.41
Substation #4	2°33'36.19"N, 102°55'50.33"E	12.41	0.3
Juaseh River			
Substation #1	2°28'54.73"N, 103°5'46.44"E	8.53	0.43
Substation #2	2°27'2.69"N, 103°4'53.80"E	2.2	0.51
Substation #3	2°28'4.56"N, 103°2'28.65"E	10.44	0.32
Substation #4	2°29'0.38"N, 102°58'37.80"E	12.87	0.45
Labis River			
Substation #1	2°25'42.55"N, 103°4'49.99"E	5.96	0.2
Substation #2	2°25'20.63"N, 103°3'54.39"E	4.63	0.05
Substation #3	2°25'7.86"N, 103°3'24.16"E	2.77	0.27
Substation #4	2°24'37.16"N, 103°3'3.55"E	7.97	0.15

merged vegetation along the river, and any potential area within the river that was unreachable by the electro-fisher. Both approaches were conducted with an hour’s effort of fishing. All the substations sampled its freshwater fish once throughout the sampling session.

The collected fish were temporarily kept in a pail filled with water before being brought back for examination. When possible, the fish were identified to species level and released back into the river. Those that were unpreserved in 10% formalin solution were taken back to the laboratory for examination. Apart from a voucher for identification, additional samples were collected, and tissues from the specimens were excised for molecular analysis. Identification was conducted according to Zakaria-Ismail et al. (2020). All fish specimens were deposited in the Zoological Collection of Forest Research Institute Malaysia (**FRIM**).

Statistical analysis

All the data recorded in the field were transferred into a master list in the Microsoft Excel software with additional information on the taxonomic order. Then, the data were sorted into respective substations, and specimens of each species were counted. Also, data on water quality at each substation were analyzed and presented in Suppl. material 1.

We defined the relative abundance as (RA), $RA = (\text{Number of individuals for species A} / \text{Total number of individuals}) \times 100\%$. Species diversity can be defined as the species richness in a certain area during a certain period. Hence, we constructed a comparison boxplot to show the differences in species richness among the investigated rivers.

In order to investigate the differences in fish composition between substations, we combined fish data from each substation into four different groups; namely group 1 which consists of all substations #1, group 2 with substations #2; group 3 comprised of substations #3 and group 4 with all substations #4. Analysis of non-metric multi-dimensional scaling (NMDS) was conducted to infer the fish composition using the metaMDS function in the R package vegan 2.5–3 (R Core Team, Vienna, Austria). We conducted a permutational analysis of variance (PERMANOVA) to compare the fish assemblages at each substation. Before PERMANOVA, the multivariate homogeneity

of group dispersions was tested using the function *betadisper*, and there was no difference in dispersion between groups/substations ($F = 0.63$, $p = 0.61$).

Results

Fish assemblage composition

In total, sixty-six freshwater fish species, including 841 specimens, were collected in the 12 substations of these three rivers (Juaseh, Labis, and Segamat) flowing off from the western part of the ERL, distributed in 43 genera, 18 families and six orders (Table 2). Among them, the most species-rich order was Cypriniformes (26 genera, seven families and 39 species), followed by the order Siluriformes (six genera, four families and 10 species). Anabantiformes and Synbranchiformes with three families comprise five genera of eight and five species, respectively. Order Gobiiformes only recorded a single family and a single genus and species.

Cyprinidae was the most dominant family, with 17 species, comprising 52% of the total abundance collected from all the substations (Table 1), followed by Danionidae and Cobitidae, with six species each. The families with the lowest number of species were Pristolepididae, Belonidae, Vaillantellidae, Siluridae, Eleotridae, and Synbranchidae, with a single species each.

Crossocheilus obscurus Tan & Kottelat, 2009 was the most abundant species collected in all the substations with a total of 85 collected specimens, followed by *Barbodes sellifer* Kottelat & Lim, 2021 and *Mystacoleucus obtusirostris* (Valenciennes, 1842), with 56 and 52 specimens, respectively. *Pangio malayana* (Tweedie, 1956), *Epalzeorhynchos kalopterum* (Bleeker, 1850), *Osteochilus microcephalus* (Valenciennes, 1842), *Nemacheilus selangoricus* Duncker, 1904, *Vaillantella maassi* Weber & de Beaufort, 1912, *Parachela oxygastroides* (Bleeker, 1852), and *Leiocassis micropogon* (Bleeker, 1852) were the species with singleton sample collected throughout the sampling session.

Fish composition among rivers and substations

Among the three rivers investigated, Segamat River recorded the highest species richness with 45 species, followed by Juaseh River with 36 species, and Labis River with 34 species of freshwater fishes. Among the investigated rivers, based on standardized effort, the highest number of specimens were captured in Segamat River (355 specimens), followed by Labis River (262 specimens), and the least was in Juaseh River (224 specimens) (Fig. 2).

The fish composition among the substations varied significantly. Substation #1 recorded 33 species, as the species richness increased in substation #2 (with 40 species) and substation #3 (41 species). However, the species richness recorded at substation #4 showed a marked reduction to 29 species compared to the other substations (Table 2). The fish compositions in substation #1 of the three rivers were grouped separately

Table 2. List of freshwater fish, relative abundance and originality that were collected from the three rivers flowing off western Endau-Rompin Landscape.

Species	Sg. Juaseh				Sg. Labis				Sg. Segamat				Relative Abundance (%)	Originality	
	SJ1	SJ2	SJ3	SJ4	SL1	SL2	SL3	SL4	SG1	SG2	SG3	SG4			
Anabantiformes															
Channidae															
<i>Channa limbata</i> (Cuvier 1831)					3				2				0.59	Native	
<i>Channa lucius</i> (Cuvier 1831)	1				1	3		1					0.71	Native	
<i>Channa melasoma</i> (Bleeker, 1851)					7								0.83	Native	
<i>Channa striata</i> (Bloch, 1793)			1									1	0.24	Native	
Osphronemidae															
<i>Betta pugnax</i> (Cantor, 1849)					8	7	3						2.14	Native	
<i>Trichopodus trichopterus</i> (Pallas, 1770)			1		1				1	1			0.48	Native	
<i>Trichopsis vittata</i> (Cuvier, 1831)			4	2									0.71	Native	
Pristolepididae															
<i>Pristolepis grootii</i> (Bleeker, 1852)		2	2	1									0.59	Native	
Beloniformes															
Belonidae															
<i>Xenentodon cancila</i> (Hamilton, 1822)											1		0.12	Native	
Zenarchopteridae															
<i>Dermogenys collettei</i> Meisner, 2001			3									2	0.59	Native	
<i>Hemirhamphodon pogonognathus</i> (Bleeker, 1853)					16	5	8						3.45	Native	
Cypriniformes															
Balitoridae															
<i>Balitoropsis zollingeri</i> (Bleeker, 1853)						1		1					0.24	Native	
<i>Homaloptera ogilviei</i> Alfred, 1967									5	5	5		1.78	Native	
<i>Homaloptera parclitella</i> Tan & Ng, 2005	5	2							4				1.31	Native	
<i>Homalopteroides tweediei</i> (Herre, 1940)											6	2	0.95	Native	
<i>Pseudohomaloptera</i> (Leonardi Hora, 1941)								2					0.24	Native	
Cobitidae															
<i>Acantopsis dialuzona</i> Van Hasselt, 1823			3		1				1	8	5	1	2.26	Native	
<i>Aperioptus pictorius</i> Richardson, 1848										1	1		0.24	Native	
<i>Pangio malayana</i> (Tweedie, 1956)											1		0.12	Native	
<i>Pangio muraeniformis</i> (de Beanfort, 1933)						3							0.36	Native	
<i>Pangio piperata</i> Kottelat & Lim, 1993											2		0.24	Native	
<i>Pangio semicincta</i> (Fraser-Brunner, 1940)			19										2.26	Native	
Cyprinidae															
<i>Barbodes lateristriga</i> (Valenciennes, 1842)	2				8	8	6	1			1		3.09	Native	
<i>Barbodes sellifer</i> (Kottelat & Lim, 2021)	9	4	3	2	17	5	9	3	2		1	3	6.9	Native	
<i>Crossocheilus obscurus</i> Tan & Kottelat, 2009	3	21	1	2			2			55	1		10.11	Native	
<i>Crossochilus oblongus</i> Kahl & Van Hasselt, 1823									20				2.38	Native	
<i>Cyclocheilichthys apogon</i> (Valenciennes, 1842)			2	3									0.59	Native	
<i>Cyclocheilichthys armatus</i> (Valenciennes, 1842)								1		2			0.36	Native	
<i>Epalzeorhynchos kalopterum</i> (Bleeker, 1850)										1			0.12	Native	
<i>Hampala macrolepidota</i> Kuhl & Van Hasselt, 1823	1	1					2	3		1			0.95	Native	
<i>Labiobarbus leptocheilus</i> (Valenciennes, 1842)				12			3	2		12		6	4.16	Native	
<i>Lobocheilos rhabdoura</i> (Fowler, 1934)	8	2							7	14	2		3.92	Native	
<i>Mystacoleucus obtusirostris</i> (Valenciennes, 1842)	1	3	7	7						28	5	1	6.18	Native	
<i>Neolissochilus soroides</i> (Duncker, 1904)	6								4				1.19	Native	
<i>Osteochilus microcephalus</i> (Valenciennes, 1842)				1									0.12	Native	
<i>Osteochilus scapularis</i> Fowler, 1939									7				0.83	Native	
<i>Osteochilus vittatus</i> (Valenciennes, 1842)			2		1		7			6	3	5	2.85	Native	
<i>Osteochilus waandersii</i> (Bleeker, 1853)	3			4	7	2	3	8	9	11		2	5.83	Native	
<i>Poropuntius normani</i> Smith, 1931					5	6			11				2.62	Native	
Danionidae															
<i>Luciosoma setigerum</i> (Valenciennes, 1842)											3		0.36	Native	
<i>Rasbora bankanensis</i> (Bleeker, 1853)					10								1.19	Native	
<i>Rasbora elegans</i> Volz, 1903	6	5	11	2	11	6	6			1			5.71	Native	

Species	Sg. Juaseh				Sg. Labis				Sg. Segamat				Relative Abundance (%)	Originality
	SJ1	SJ2	SJ3	SJ4	SL1	SL2	SL3	SL4	SG1	SG2	SG3	SG4		
<i>Rasbora myersi</i> Brittan, 1954				3			1		1	2			0.83	Native
<i>Rasbora paucisqualis</i> Ahl, 1935						7			8		8		1.78	Native
<i>Trigonostigma heteromorpha</i> (Duncker, 1904)					12								1.43	Native
Nemacheilidae														
<i>Nemacheilus paucimaculatus</i> Bohlen & Šlechtová, 2011					1	1	1	3		11			2.02	Native
<i>Nemacheilus selangoricus</i> Duncker, 1904											1		0.12	Native
Vaillantellidae														
<i>Vaillantella maassi</i> Weber & de Beaufort, 1912										1			0.12	Native
Xenocyprididae														
<i>Parachela oxygastroides</i> (Bleeker, 1852)			1										0.12	Native
<i>Oxygaster anomalura</i> Van Hasselt, 1823											2		0.24	Native
Gobiiformes														
Eleotridae														
<i>Oxyeleotris marmorata</i> (Bleeker, 1852)			4										0.48	Native
Siluriformes														
Bagridae														
<i>Leiocassis micropogon</i> (Bleeker, 1852)										1			0.12	Native
<i>Mystus castaneus</i> Ng, 2002	2				2				2				0.71	Native
<i>Mystus singaringan</i> (Bleeker, 1846)		1	4										0.59	Native
<i>Pseudomystus leiacanthus</i> (Weber & de Beaufort, 1912)					1					2			0.36	Native
Clariidae														
<i>Clarias batrachus</i> (Linnaeus, 1758)			2			1							0.36	Native
<i>Clarias leiacanthus</i> Bleeker, 1851					3	2							0.59	Native
Siluridae														
<i>Silurichthys hasseltii</i> Bleeker 1858					2	1			1				0.48	Native
Sisoridae														
<i>Glyptothorax fuscus</i> Fowler, 1934	1							1	2	9		5	2.14	Native
<i>Glyptothorax platypogonoides</i> (Bleeker, 1855)						3					3		0.71	Native
<i>Glyptothorax schmidti</i> (Volz, 1904)	1				1								0.24	Native
Synbranchiformes														
Syngnathidae														
<i>Doryichthys martensii</i> (Peters, 1868)			9								4	2	1.78	Native
<i>Doryichthys deokhatoides</i> (Bleeker, 1854)				1						4	4		1.07	Native
Synbranchidae														
<i>Monopterus javanensis</i> Lacepede, 1800			1			2	3	1					0.83	Native
Mastacembelidae														
<i>Macrognathus maculatus</i> (Cuvier, 1832)	1								1	1		1	0.48	Native
<i>Mastacembelus favius</i> Hora 1924		11	1	1									1.55	Native

compared with other substations, as shown in the NMDS ordination (Fig. 3). The fish structure in substations #2, #3, and #4 overlapped.

Discussions

The fish assemblage presented in this study is the first insight into fish diversity found in the Juaseh, Segamat, and Labis Rivers, flowing off from the western part of the ERL. A total of 66 freshwater fish species were recorded in the three rivers, representing approximately 10% of freshwater fish diversity found in Malaysian territory. Fishes of the order Cypriniformes are the main freshwater inhabitants and contribute to almost




Figure 2. The boxplot shows the species richness of freshwater fish for all three investigated rivers in the western part of the Endau-Rompin Landscape.




Figure 3. NMDS analysis for each substation examined along study areas. The composition of fish on substations #1 differs significantly from substations of #2, #3 and #4. Each convex hull represents Group 1 = substations #1, Group 2 = Substations 2, Group 3 = Substations #3 and Group 4 = Substations #4.

10% of all teleost species globally (Wood 2019). Most tropical freshwater fishes in Asia are dominated by Cypriniformes (barbs and loaches), Siluriformes (catfishes), anabantoids (gouramis, climbing perches, and snakeheads), and mastacembelids (spiny eels). In addition, several freshwater species come from estuaries or marine environments, such as: Beloniformes (halfbeaks and needlefish), Atheriniformes (silversides), Eleotridae (sleeper gobies), Gobiidae (goby), and Tetraodontidae (pufferfish) (Wine-miller et al. 2008). Across the investigated rivers, the family Cyprinidae was dominant and accounted for almost half of the total catch. Dominance by the family Cyprinidae in tropical rivers is common due to their highly morphological adaptations (Ward-Campbell et al. 2005), wide distribution (Mohsin and Ambak 1983), and is supported by previous studies (Kaviarasu et al. 2013; Rashid et al. 2015; Ng et al. 2019, Soo et al. 2021) conducted in Peninsular Malaysia and Borneo.

Families, such as Belonidae, Vaillantellidae, and Eleotridae, were less sampled in this study. A similar result was reported in the Keniyam River, in National Park Taman Negara Pahang by Mohd-Azham and Singh (2019), with a low abundance of belonid species. The family Siluridae (silurid catfishes) have 17 species recorded in Peninsular Malaysia; however, this study only managed to capture four specimens of the species *Silurichthys hasseltii* Bleeker, 1858. Based on the documentation of freshwater fishes in Peninsular Malaysia by Zakaria-Ismail et al. (2019), most silurid species are rare and restricted to particular rivers in Peninsular Malaysia. This justifies the fact that species of this family usually present low abundance, and the chances of collecting species of silurid is extremely low.

Crossocheilus obscurus of the family Cyprinidae was the most abundant species among all three rivers in this study. This species was described by Tan and Kottelat (2009) from the middle of the Batang Hari drainage of Sumatra. The species has also been discovered in Malaysia, specifically in the Muar River of Johor and the Kampar River in Perak. Even though Zakaria-Ismail et al. (2019) confirmed that the species is uncommon in Peninsular Malaysia, our study was able to gather a sizable number of *C. obscurus* specimens from the Segamat and Juaseh Rivers (total specimens of 85). It could be a spawning ground for *C. obscurus* in the Juaseh and Segamat Rivers, as the abundance was high, and the species dominated several substations along both rivers. Also, habitat preferences for *C. obscurus* are related to fast-flowing streams with clear water and the bottom substrates of rocks and submerged logs. Such conditions were observed in the substations where the species were recorded, providing an optimum condition for its persistency. Conversely, most species that were collected in this study are low in abundance. However, these fishes were commonly present throughout Peninsular Malaysia (Zakaria-Ismail et al. 2019).

Fish species composition is marked by habitat heterogeneity, environmental gradients, and human activity (Sehr and Keckeis 2017; Shukla and Bhat 2017) at local and regional scales (Benke et al. 2011; Hasegawa et al. 2017; Nicol et al. 2017; Cheng et al. 2019). Comparably, the fish richness in this study increases across the substations following the river's gradient. Basically, substation #1 is located at the headwater of each river, whereas substations #2 and #3 are a feature of midstream, and substation

#4 is downstream. The fish assemblage reflects the gradual changes in structural and physicochemical properties from the upper to the lower reaches (Petry and Schulz 2006). A higher species richness was expected in substation #4 as the fish species richness progressed toward downstream (Horwitz 1978; Vannote et al. 1980), but surprisingly, fish assemblage was much lower. Such findings may contribute to land-use change and anthropogenic disturbance downstream (Benejam et al. 2016; Leitão et al. 2018). It was observed that the intensity of human settlement and agricultural activity downstream were significant compared to the upper reaches. Hence, the downstream fish composition was majorly dominated by highly tolerant species, and those species that had acquired cleaner and pristine habitats were absent. Such a suggestion was supported by the NMDS ordination result of this study. The convex hull grouped substations according to the fish composition, which resulted in substation #1 being separated from the other substations. Such results indicated that different species were present exclusively in substation #1 and absent from other substations.

Conservation of fish in the ERL and recommendations

The community's major economy in the study area is oil palm plantations and agricultural estates. The current land-use changes in the study areas, which humans mainly cause, would directly impact negatively on freshwater fish communities. It was suggested that any form of logging activity, including selective logging, widely practiced by foresters, could create a positive ecosystem for freshwater fish (Iwata et al. 2003; Wilkinson et al. 2018). Although there was no significant difference in fish diversity between logged forest and oil palm catchments in the Sabah River reported by Wilkinson et al. (2018), the forests can be constructive and influence the diversity of freshwater fish communities by increasing the heterogeneity in physical structures and food materials in freshwater habitats (Lo et al. 2020). The anthropogenic land cover change could also serve to increase the sedimentation, changes in water pH, and the availability of food materials that directly influence the number of aquatic organisms (Lo et al. 2020). Consequently, the understanding of habitat alteration, especially between forest and aquatic ecosystems, should be enhanced to provide a systematic framework for conservation programs.

Disturbances in the riparian zone along the rivers could have harmful impacts on the terrestrial communities and indirectly affect the freshwater ecosystem (Grimm et al. 2008; Vörösmarty et al. 2010). Riparian zones are the landscapes that form an interface between terrestrial and aquatic ecosystems, with disproportionate influences on the dynamics of the food web and the function of the ecosystem in both habitats. The plant communities along the riverbanks and margins are known as riparian vegetation. They are very significant for fish communities, as they influence light regimes, water quality, thermal dynamics, habitat, and food availability. The riparian buffer along Juaseh, Segamat, and Labis rivers, especially downstream, was not clearly visible as most zones were destroyed or encroached by human activities. The loss of riparian vegetation situated along these rivers could eradicate the food sources and breeding

areas of the fish communities. A successful idea of adapting buffer zones has been reviewed in India, where the program, Freshwater Fish Safe Zone, has effectively aided in freshwater fish conservation (Gupta et al. 2014). Therefore, safeguarding the network of the river-riparian zone would be essential in rehabilitating the degraded habitats, which resulted from the urbanization (Coelho et al. 2014).

Freshwater fishes are important sources of protein to the rural communities that live in the inland and riverine rural areas. However, many freshwater fishes in Malaysia have been overharvested, especially the large-sized and game fishes, including Mahseer and *Probarbus jullieni* Sauvage, 1880 (Chong et al. 2010). Overharvesting mostly occurred in habitats that lack surveillance, which could lead to the decreasing number of freshwater fishes or extinction of the species in the wild. For certain species, such as Betta, it is believed to be overharvested due to trading for the ornamental industry since Betta is very famous and has always been a popular preference among ornamental fish enthusiasts because of its color and shape in Malaysia. Therefore, relevant authorities should take proactive measures in curbing overharvesting by enforcing, monitoring, and guarding the natural habitats besides introducing seasonal harvesting, which could assist in maintaining the wild population of freshwater fish.

The establishment of policies or management plans for the freshwater ecosystem is a matter of requirement, but enforcement could be more crucial. Even though several conservation policies have been put in place in Malaysia for many years, the government continues to poorly manage most of the ecosystems due to the lack of stakeholder participation, inefficient conservation programs, and failure to raise community awareness (Abdullah et al. 2014). Engagement by pertinent stakeholders and authorities should be more holistic and covered by all layers of administration. In this study, the relevant authorities responsible for freshwater fish and ecosystem rely on the Department of Fisheries, State Forestry Department, and the district official since the rivers transcend multi-jurisdictions. These authorities should be advised based on scientific facts to help them design, develop, and execute any rehabilitation effort within their jurisdiction. The involvement of local communities is crucial in ensuring that any conservation and management plan in place is successful. According to Rampheri and Dube (2020), local communities play an essential role in executing management plans since most locals have extensive knowledge and experience in managing and conserving natural resources while safeguarding their livelihoods. The local community should be further informed on nature and its benefits that will likely influence their lives and improve their awareness of conserving the surrounding ecosystems. The scientific society in Malaysia should be more engaging in conserving the freshwater ecosystem by conducting in-depth research, especially on long-term monitoring. The relationship between species and the ecosystem is complex, and the diversity of monitoring requirements must be considered and combined (Deines et al. 2017; Allison and Mills 2018). The conservation measures should be evidence-based to implement the best strategy to protect and rehabilitate the existing ecosystems within the country.

Conclusion

The knowledge presented here is the first documentation on the freshwater fish from the Juaseh, Labis, and Segamat rivers in Malaysia. The diversity and composition of fish species in these rivers differed suggestively, and the distribution along the river gradient changes gradually from the headwater to downstream. The information would serve as a baseline information for key authorities and stakeholders to preserve and conserve the biodiversity inhabiting freshwater ecosystems in Malaysia, in a sustainable management manner.

Acknowledgments

This project is funded by the Ministry of Higher Education Malaysia (MOHE) under the Fundamental Research Grant Scheme FRGS/1/2020/WAB11/KATS//1 (UTHM-K353) and UTHM-GPPS-H702 postgraduate grant by Universiti Tun Hussein Onn Malaysia. Most of the equipment used during the investigation was supported by project RMk-12 P23085100210003. The authors would like to thank Dato' Indera Mohd Ridza Bin Awang, Director General of Forestry Department of Peninsular Malaysia (JPSM), and Director of Johor State Forestry Department for the permit given to assess the sites within the permanent forest reserve of Labis (Permit No: PHDJU 37/2021; PHDJU 104/2022); and we would like to thank Dato' Abdul Kadir Abu Hashim, Director General of Department of Wildlife and National Park (DWNP) Peninsular Malaysia for the research permit (B-00381-15-22). The authors would also like to thank Tuan Haji Zainudin bin Haji Abd Wahab, Director of the Department of Fishery of Johor, for their permission and assistance during data collection and sampling. Special mentions also go to Wendy Yong Sze Yee for generating the study site map and Dr. Lillian Chua, Director of Forest Biodiversity Division of Forest Research Institute Malaysia (FRIM), for her kind guidance and advice. The authors acknowledge MOHE, FRIM, and Universiti Tun Hussein Onn Malaysia (UTHM) for providing the necessary funding, facilities, and assistance. Lastly, the authors thank Enago (www.enago.com) for their English language review and editing services.

References

- Abdullah K, Said AM, Omar D (2014) Community-based conservation in Managing Mangrove Rehabilitation in Perak and Selangor. *Procedia: Social and Behavioral Sciences* 153: 121–131. <https://doi.org/10.1016/j.sbspro.2014.10.047>
- Allison EH, Mills DJ (2018) Counting the fish eaten rather than the fish caught. *Proceedings of the National Academy of Sciences of the United States of America* 115(29): 7459–7461. <https://doi.org/10.1073/pnas.1808755115>

- Aqilah AAR, Maryati M, Syahrir-Azrin S (2018) Diversity and species composition of butterflies (Lepidoptera: Papilionoidea) in Taka Melor Amenity Forest, Labis Forest Reserve, Segamat, Johor, Malaysia. *Serangga* 24(2): 72–80.
- Benejam L, Teixeira-de Mello F, Meerhoff M, Loureiro M, Jeppesen E, Brucet S (2016) Assessing effects of change in land use on size-related variables of fish in subtropical streams. *Canadian Journal of Fisheries and Aquatic Sciences* 73(4): 547–556. <https://doi.org/10.1139/cjfas-2015-0025>
- Benke M, Brändle M, Albrecht C, Wilke T (2011) Patterns of freshwater biodiversity in Europe: Lessons from the spring snail genus *Bythinella*. *Journal of Biogeography* 38(10): 2021–2032. <https://doi.org/10.1111/j.1365-2699.2011.02527.x>
- Chan NW (2002) Pembangunan, Pemandaran dan Peningkatan Bahaya dan Bencana Air di Malaysia: Isu, Pengurusan dan Cabaran. Penerbit Universiti Sains Malaysia.
- Chan NW (2012) Managing urban rivers and water quality in Malaysia for sustainable water resources. *International Journal of Water Resources Development* 28(2): 343–354. <https://doi.org/10.1080/07900627.2012.668643>
- Cheng D, Zhao X, Song J, Sun H, Wang S, Bai H, Li Q (2019) Quantifying the distribution and diversity of fish species along elevational gradients in the Weihe River Basin, Northwest China. *Sustainability (Basel)* 11(21): 6177. <https://doi.org/10.3390/su11216177>
- Chong VC, Lee PK, Lau CM (2010) Diversity, extinction risk and conservation of Malaysian fishes. *Journal of Fish Biology* 76(9): 2009–2066. <https://doi.org/10.1111/j.1095-8649.2010.02685.x>
- Coelho D, Hughes SJ, Varandas S, Cortes RMV (2014) Conservation benefits of riparian buffers in urban areas: The case of the Rio Corgo (north Portugal). *Fundamental and Applied Limnology* 185(1): 55–70. <https://doi.org/10.1127/fal/2014/0589>
- Darwall W, Bremerich V, De Wever A, Dell AI, Freyhof J, Gessner MO, Grossart HP, Harrison I, Irvine K, Jähnig SC, Jeschke JM, Lee JJ, Lu C, Lewandowska AM, Monaghan MT, Nejstgaard JC, Patricio H, Schmidt-Kloiber A, Stuart SN, Thieme M, Tockner K, Turak E, Weyl O (2018) The Alliance for Freshwater Life: A global call to unite efforts for freshwater biodiversity science and conservation. *Aquatic Conservation* 28(4): 1015–1022. <https://doi.org/10.1002/aqc.2958>
- Deines AM, Bunnell DB, Rogers MW, Bennion D, Woelmer W, Sayers MJ, Grimm AG, Shuchman RA, Raymer ZB, Brooks CN, Mychek-Londer JG, Taylor W, Beard Jr TD (2017) The contribution of lakes to global inland fisheries harvest. *Frontiers in Ecology and the Environment* 15(6): 293–298. <https://doi.org/10.1002/fee.1503>
- Dudgeon D, Arthington AH, Gessner MO, Kawabat Z, Knowler DJ, Leveque C, Naiman RJ, Prieur-Richard A, Soto D, Stiassny MLJ, Sullivan CA (2006) Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews of the Cambridge Philosophical Society* 81(2): 163–182. <https://doi.org/10.1017/S1464793105006950>
- Gardner RC, Finlayson M (2018) Global wetland outlook: State of the world's wetlands and their services to people. Secretariat of the Ramsar Convention.
- Gleick PH (1996) Basic water requirements for human activities: Meeting basic needs. *Water International* 21(2): 83–92. <https://doi.org/10.1080/02508069608686494>

- Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM (2008) Global change and the ecology of cities. *Science* 319(5864): 756–760. <https://doi.org/10.1126/science.1150195>
- Grooten M, Almond RE (2018) Living planet report-2018: aiming higher. WWF international.
- Gupta N, Nautiyal P, Borgohan A, Sivakumar K, Mathur VB, Chadwick MA (2014) Catch-and-release angling as a management tool for freshwater fish conservation in India. *Oryx* 50(2): 1–7. <https://doi.org/10.1017/S0030605314000787>
- Harrison I, Abell R, Darwall W, Thieme ML, Tickner D, Timboe I (2018) The freshwater biodiversity crisis. *Science* 362(6421): 1369. <https://doi.org/10.1126/science.aav9242>
- Hasegawa K, Mori T, Yamazaki C (2017) Density-dependent effects of non-native brown trout *Salmo trutta* on the species-area relationship in stream fish assemblages. *Journal of Fish Biology* 90(1): 370–383. <https://doi.org/10.1111/jfb.13185>
- Horwitz RJ (1978) Temporal variability patterns and the distributional patterns of stream fishes. *Ecological Monographs* 48(3): 307–321. <https://doi.org/10.2307/2937233>
- Iwata T, Nakano S, Inoue M (2003) Impacts of past riparian deforestation on stream communities in a tropical rainforest in Borneo. *Ecological Applications* 13(2): 461–473. [https://doi.org/10.1890/1051-0761\(2003\)013\[0461:IOPRDO\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0461:IOPRDO]2.0.CO;2)
- Kaviarasu M, Amin MHN, Rizal MR, Shahfiz MA (2013) Diversity and Distribution of Freshwater Fish of Temengor Lake, West Malaysia. In: Latif MA, Baskaran DK, Zainom K, Zulfadlan AK, Nurul II, Eza FRN, Alizah S, Syaiful MM, Najmi MH, Ahmad Ridzuan YA (Eds) *Proceedings of the 2nd Temengor Scientific Expedition, Pulau Banding (Malaysia) August 2013*, Pulau Banding Foundation, 321–334.
- Leitão RP, Zuanon J, Mouillot D, Leal CG, Hughes RM, Kaufmann PR, Villéger S, Pompeu PS, Kasper D, de Paula FR, Ferraz SFB, Gardner TA (2018) Disentangling the pathways of land use impacts on the functional structure of fish assemblages in Amazon streams. *Ecography* 41(1): 219–232. <https://doi.org/10.1111/ecog.02845>
- Lo M, Reed J, Castello L, Steel EA, Frimpong EA, Ickowitz A (2020) The influence of Forests in Freshwater Fish in the Tropics: A systematic review. *Bioscience* 70(5): 404–414. <https://doi.org/10.1093/biosci/biaa021>
- Miller EC (2021) Comparing diversification rates in lakes, rivers, and the sea. *Evolution; International Journal of Organic Evolution* 75(8): 2055–2073. <https://doi.org/10.1111/evo.14295>
- Mohsin AKM, Ambak MA (1983) *Freshwater fishes of Peninsular Malaysia*. Selangor, Malaysia. Universiti Pertanian Malaysia.
- Mohd-Azham MY, Singh HR (2019) Freshwater fish diversity and their distribution along the Keniyam River, Taman Negara Pahang, Malaysia. *IOP Conference Series: Earth and Environmental Science*, Jul 2019. IOP Publishing 269: e012034. <https://doi.org/10.1088/1755-1315/269/1/012034>
- Naiman RJ, Bilby RE (1998) *River Ecology and Management: Lessons from the Pacific Coastal Ecoregion*. Springer-Verlag, New York. <https://doi.org/10.1007/978-1-4612-1652-0>
- Ng CK, Aun-Chuan OP, Wong WL, Khoo G (2017) An overview of the status, trends and challenges of freshwater fish research and conservation in Malaysia. *Journal of Survey Fisheries Sciences*. 3(2): 7–21. <https://doi.org/10.18331/SFS2017.3.2.2>

- Ng CK, Lim TY, Ahmad AB, Khaironizam MZ (2019) Provisional checklist of freshwater fish diversity and distribution in Perak, Malaysia and some latest taxonomic concerns. *Zootaxa* 4567(3): 515–545. <https://doi.org/10.11646/zootaxa.4567.3.5>
- Nicol E, Stevens JR, Jobling S (2017) Riverine fish diversity varies according to geographical isolation and land use modification. *Ecology and Evolution* 7(19): 7872–7883. <https://doi.org/10.1002/ece3.3237>
- Petry AC, Schulz UH (2006) Longitudinal changes and indicator species of the fish fauna in the subtropical Sinos River, Brazil. *Journal of Fish Biology* 69(1): 272–290. <https://doi.org/10.1111/j.1095-8649.2006.01110.x>
- Rampheri MB, Dube T (2020) Local community involvement in nature conservation under the auspices of community-based natural resource management: A state of the art review. *African Journal of Ecology* 59(4): 799–808. <https://doi.org/10.1111/aje.12801>
- Rashid ZA, Asmuni M, Amal MNA (2015) Fish diversity of Tembeling and Pahang Rivers, Pahang, Malaysia. *Check List* 11(5): 1753–1753. <https://doi.org/10.15560/11.5.1753>
- Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PT, Kidd KA, Tyson J, MacCormack TJ, Olden JD, Ormerod SJ, John P, Smol JP, Taylor WW, Klement TK, Vermaire JC, Dudgeon D, Cooke SJ (2019) Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews of the Cambridge Philosophical Society* 94(3): 849–873. <https://doi.org/10.1111/brv.12480>
- Saaban S, Yasak MN, Gumal M, Oziar A, Cheong F, Shaari Z, Tyson M, Hedges S (2020) Viability and management of the Asian elephant (*Elephas maximus*) population in the Endau Rompin landscape, Peninsular Malaysia. *PeerJ* 24(8): e8209. <https://doi.org/10.7717/peerj.8209>
- Sala OE, Chapin 3rd FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A, Leemans R, Lodge DM, Mooney HA, Oesterheld M, Poff NL, Sykes MT, Walker BH, Walker M, Wall DH (2000) Global biodiversity scenarios for the year 2100. *Science* 287(5459): 1770–1774. <https://doi.org/10.1126/science.287.5459.1770>
- Sehr M, Keckeis H (2017) Habitat use of the European mudminnow *Umbra krameri* and association with other fish species in a disconnected Danube side arm. *Journal of Fish Biology* 91(4): 1072–1093. <https://doi.org/10.1111/jfb.13402>
- Shukla R, Bhat A (2017) Environmental drivers of α -diversity patterns in monsoonal tropical stream fish assemblages: A case study from tributaries of Narmada basin, India. *Environmental Biology of Fishes* 100(7): 749–761. <https://doi.org/10.1007/s10641-017-0601-6>
- Soo CL, Nyanti L, Idris NE, Ling TY, Sim SF, Grinang J, Ganyai T, Lee KS (2021) Fish biodiversity and assemblages along the altitudinal gradients of tropical mountainous forest streams. *Scientific Reports* 11(1): e16922. <https://doi.org/10.1038/s41598-021-96253-3>
- Tan HH, Kottelat M (2009) The fishes of the Batang Hari drainage, Sumatra, with description of six new species. *Ichthyological Exploration of Freshwaters* 20(1): 13–69.
- Tickner D, Opperman JJ, Abell R, Acreman M, Arthington AH, Bunn SE, Cooke SJ, Dalton J, Darwall W, Edwards G, Harrison I, Hughes K, Jones T, Leclère D, Lynch AJ, Leonard P, McClain ME, Muruven D, Olden JD, Ormerod SJ, Robinson J, Tharme RE, Thieme M, Tockner K, Wright M, Young L (2020) Bending the Curve of Global Freshwater Biodiversity Loss. *Bioscience* 70(4): 330–342. <https://doi.org/10.1093/biosci/biaa002>

- Val P, Lyons NJ, Gasparini N, Willenbring JK, Albert JS (2022) Landscape evolution as a diversification driver in freshwater fishes. *Frontiers in Ecology and Evolution* 9: e788328. <https://doi.org/10.3389/fevo.2021.788328>
- Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980) The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37(1): 130–137. <https://doi.org/10.1139/f80-017>
- Vörösmarty CJ, McIntyre PB, Gessner MO, Dudgeon D, Prusevich A, Green P, Glidden S, Bunn SE, Sullivan CA, Liermann CR, Davies PM (2010) Global threats to human water security and river biodiversity. *Nature* 467(7315): 555–561. <https://doi.org/10.1038/nature09440>
- Ward-Campbell BMS, Beamish FWH, Kongchaiya C (2005) Morphological characteristics in relation to diet in five coexisting Thai fish species. *Journal of Fish Biology* 67(5): 1266–1279. <https://doi.org/10.1111/j.1095-8649.2005.00821.x>
- Wilkinson CL, Yeo DCJ, Tan HH, Fikri AH, Ewers RM (2018) The availability of freshwater fish resources is maintained across a land-use gradient in Sabah, Borneo. *Aquatic Conservation* 28(5): 1044–1054. <https://doi.org/10.1002/aqc.2920>
- Winemiller KO, Agostinho AA, Érica Pellegrini Caramaschi EP (2008) Fish Ecology in Tropical Streams. In: David Dudgeon D (Ed.) *Tropical Stream Ecology*. Academic Press, 107–111. <https://doi.org/10.1016/B978-012088449-0.50007-8>
- Wood CM (2019) Internal spatial and temporal CO₂ dynamics: Fasting, feeding, drinking, and the alkaline tide. In: Grosell M, Munday, PL, Farrell AP, Brauner CJ (Eds) *Fish Physiology: Carbon Dioxide* Elsevier Inc., 37: 245–286. <https://doi.org/10.1016/bs.fp.2019.07.003>
- Zakaria-Ismail M, Fatimah A, Khaironizam MZ (2019) *Fishes of the freshwater ecosystems of Peninsular Malaysia*. LAP Lambert Academic Publishing, 356 pp.

Supplementary material I

Eleven environmental variables measured in each substation of rivers Juaseh, Segamat, and Labis of ERL.

Authors: Munian Kaviarasu, Farah Farhana Ramli, Lokman Mohd Ilham Norhakim, Nursyuhada Othman, Nur Aina Amira Mahyuddin, Hidayah Haris, Nur Hartini Sariyati, Mohd Faudzir Najmuddin, Salim Aman, Salman Faris Zaharin, Muhammad Abu Bakar Abdul-Latiff

Data type: Environmental data.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/natureconservation.50.86090.suppl1>